

TABLE IV - FISH OCCURRENCE IN ANACONDA-PINTLER LAKES

LAKE NAME	DRAINAGE ¹	SPECIES ²	NATURAL REPRODUCTION	AMT. USE	COMMENTS
Alpine Lakes	Alpine Ck., Brt	Fishless	NA	Low	
"Annie", (T3 N R16 W Sec 35)	M Fk. Rk Ck	Fishless		Med	Natural springs
Bear	Beaver Ck., BH	Fishless	?	Low	Shallow
Buck	Buck Ck., Brt	Fishless	NA	Low	Shallow
Carmine	Carmine Ck., Brt	Fishless	NA	Low	
Carpp	Carpp Ck., RkCk	WCT	?	High	High day use, poor nutrition, possible winterkill
Lower Carpp	Carpp Ck., RkCk			High	
Upper Carpp	Carpp Ck, RkCk	Fishless		Med	Shallow
Continental	Thompson Ck., BH	WCT	?	Low	Shallow, formerly stocked with RB, 1946
Crystal	Thompson Ck., BH	RB	yes	Med.	
Edith	Falls Fk., RkCk	RB, GR, WCT	yes	High	"Natural reproduction provides more recruitment than desirable for maximum growth"
Emerald	EFk.LaMarche, BH	Fishless	NA	Low	
Flower	Page Ck., RkCk.	Fishless	NA	Med.	Shallow
Hicks	WFk.LaMarche, BH	RB	yes?	Med	
Hidden	Brt	WCT	yes	High	
Hope	Hope Ck., Brt	WCT	yes	Med	Increasing Use
Ivanhoe	M.Fk. RkCk	RB	yes	Med	
Johnson	Falls Fk., RkCk	WCT, LNS	yes	High	Most heavily used area in A-P;
Kelly	EFk Brt	Fishless		High	Shallow
LaMarche	LaMarche, BH			Low	
Lion	Thompson Ck., BH	RB	yes	Med	
"Little Annie" T3N R15W Sec 16	Tamarack Ck, RkCk			Med	

¹The Bitterroot and Rock Creek drainages are west of the Continental Divide. Both drainages have native Westslope Cutthroat and Bull Trout populations. The Big Hole drainage is east of the Continental Divide and has Upper Missouri Westslope Cutthroat.

² WCT=Westslope Cutthroat; RB=Rainbow; GR=Non-Native Grayling; LNS=Long-Nosed Sucker

FISHERIES MANAGEMENT UPDATE

Little Johnson	M Fk RkCk			Med	
Lost Lakes	EFk Fish Trap, BH	Fishless		Low	
Martin	Falls Fk, RkCk	Fishless		Low	Shallow

TABLE IV - FISH OCCURRENCE IN ANACONDA-PINTLER LAKES (continued)

LAKE NAME	DRAINAGE ³	SPECIES ⁴	NATURAL REPRODUCTION	AMT. USE	COMMENTS
Mosquito	Thompson Ck, BH				
Mystic	Howell Ck., BH	RB	yes	Med	
Oreamnos	PintlerCk., BH	RB	yes	High	Fish freeze out; repeatedly stocked in past
Page	Page Ck., RkCk	Fishless?		Med	Historically planted with RB and CT
Park Lakes	Park Ck, EFk Brt	Fishless		Low	
Phlox	Hell Roaring, BH	Fishless		Low	Shallow
Phyllis	M Fk Rk Ck	RB, CT	?	Med	
Upper Phyllis	RkCk	CT	?	High	
Queener Pond (Horse Camp)	Seymour, BH			Med	
Rainbow	BH	RB	yes	High	
Ripple	EFk Btr			High	
Sauer	Sauer Ck., RkCk	?		Low	Historically planted with CT
Sawed Cabin	Pintler Ck., BH	Fishless		Med	Shallow
Spruce	Spruce Ck, RkCk				
Surprise	Hell Roaring, BH	Fishless		Med	Shallow
Tamarack	Carp Cr, RkCk	Planted WCT	no	High	Increased use since stocked in 1993
Unnamed, West, Warren Lake	BH	Fishless		Low	
Unnamed, below Queener Mnt.	Seymour Ck, BH	Fishless		Low	
Unnamed lake T3N, R15W, Sec 21, NW1/4	Carp Cr., RkCk.	RB	yes	Low	

³The Bitterroot and Rock Creek drainages are west of the Continental Divide. Both drainages have native Westslope Cutthroat and Bull Trout populations. The Big Hole drainage is east of the Continental Divide and has an East slope strain of Westslope Cutthroat.

⁴ WCT=Westslope Cutthroat; RB=Rainbow; GR=Non-Native Grayling; LNS=Long-Nosed Sucker

FISHERIES MANAGEMENT UPDATE

Upper Seymour	Seymour Ck, BH	RB	yes	High	
Violet	Hell Roaring, BH				
Warren	LaMarche, BH	RB	yes	Med.	Stocked in past